



NEWS FROM THE UNIVERSITY OF TASMANIA, AUSTRALIA

Media Release

Chiefs of Staff, News Directors

Wednesday 11 April 2018

Hotter, longer, more frequent – global marine heatwaves on the rise

An [international study in *Nature Communications*](#) co-authored by researchers from the ARC Centre of Excellence for Climate Extremes (CLEX) and IMAS reveals globally marine heatwaves have increased over the past century in number, length and intensity as a direct result of warming oceans.

From 1925-2016, the study found the frequency of marine heatwaves had increased on average by 34% and the length of each heatwave had increased by 17%. Combined, this led to a 54% increase in the number of marine heatwave days every year.

“Our research also found that from 1982 there was a noticeable acceleration of the trend in marine heatwaves,” said lead author, Dr Eric Oliver from Dalhousie University, Canada.

“While some of us may enjoy the warmer waters when we go swimming, these heatwaves have significant impacts on ecosystems, biodiversity, fisheries, tourism and aquaculture. There are often profound economic consequences that go hand in hand with these events.”

Some recent examples show just how significant marine heatwave events can be.

- In 2011, Western Australia saw a marine heatwave that shifted ecosystems from being dominated by kelp to being dominated by other seaweed. That shift remained even after water temperatures returned to normal;
- In 2012, [a marine heatwave in the Gulf of Maine](#) led to an increase in lobsters but a crash in prices that seriously hurt the industry’s profits;
- Persistent warm water in the north Pacific from 2014-2016 led to fishery closures, mass strandings of marine mammals and harmful algal blooms along coastlines. That heatwave even changed large-scale weather patterns in the Pacific Northwest;
- More recently, [Tasmania’s intense marine heatwave in 2016](#) led to disease outbreaks and slowing in growth rates across aquaculture industries.

The researchers used a variety of observational datasets to reveal the trend of increasing marine heatwaves, combining satellite data with a range of century long datasets taken from ships and various land based measuring stations.

They then removed the influences of natural variability caused by the El Nino Southern Oscillation, the Pacific Decadal Oscillation and the Atlantic Multidecadal Oscillation to find the underlying trend.

“There was a clear relationship between the rise in global average sea-surface temperatures and the increase in marine heatwaves, much the same as we see increases in extreme heat events related to the increase in global average temperatures,” said IMAS co-author Professor Neil Holbrook.

“With more than 90% of the heat from human caused global warming going into our oceans, it is likely marine heatwaves will continue to increase.

“The next key stage for our research is to quantify exactly how much they may change.

“The results of these projections are likely to have significant implications for how our environment and economies adapt to this changing world,” Professor Holbrook said.

More information on marine heatwaves can be found at www.marineheatwaves.org.

Downloadable content: <https://cloudstor.aarnet.edu.au/plus/s/BVOdUdR2y0B3yzA>

- **Video grabs of Professor Neil Holbrook talking about the increase in marine heatwaves**

Media contacts:

Alvin Stone
Media and Communications Manager
ARC Centre of Excellence for Climate Extremes, and
ARC Centre of Excellence for Climate System Science
University of New South Wales
+61 2 9385 8953, +61 (0) 418 617 366
alvin.stone@unsw.edu.au

Andrew Rhodes
Communications Manager
Institute for Marine and Antarctic Studies, University of Tasmania
+61 3 6226 6683, +61 (0) 417 239 537, a.j.rhodes@utas.edu.au

Nicole Comeau
Faculty of Science - Communications
Dalhousie University, Canada
(902) 494-8443
(902) 223-2446

Information released by:

Communications and Media Office, University of Tasmania
Phone: 61 3 6226 2124
Email: Media.Office@utas.edu.au